

主な研究課題・発表代表論文

病理形態学講座（旧病理学第一講座）Pathology

研究領域 構造解析病態医学

教授 池田 栄二 Eiji Ikeda

Web ページ：<https://ds.cc.yamaguchi-u.ac.jp/~pathol1/>

主な研究課題

- ・神経系血管バリアー機能の人為的制御手法の確立に向けた基礎研究
- ・新規組織診断基準および治療法開発を目指したグリオーマの病態解析
- ・成人肺の再生戦略研究

発表代表論文

- 1) Cui D, Yamamoto K, Ikeda E: High-mannose-type glycan of basigin in endothelial cells is essential for the opening of blood-brain barrier induced by hypoxia, cyclophilin A or TNF- α . *Am J Pathol* (in press)
- 2) Nakada-Honda N, Cui D, Matsuda S, Ikeda E : Intravenous injection of cyclophilin A realizes the transient and reversible opening of barrier of neural vasculature through basigin in endothelial cells. *Sci Rep* 11:19391. doi: 10.1038/s41598-021-98163-w, 2021
- 3) Cui D, Arima M, Hirayama T, Ikeda E : Hypoxia-induced disruption of neural vascular barrier is mediated by the intracellular induction of Fe (II) ion. *Exp Cell Res* 379 : 166-171, 2019. doi : 10.1016/j.yexcr.2019.04.003. Epub 2019 Apr 7
- 4) Kimura T, Cui D, Kawano H, Yoshitomi-Sakamoto C, Takakura N, Ikeda E: Induced expression of GINS complex is an essential step for reactivation of quiescent stem-like tumor cells within the peri-necrotic niche in human glioblastoma. *J Cancer Res Clin Oncol.* 2018 Nov 21. doi : 10.1007/s00432-018-2797-z.
- 5) Arima M, Cui D, Kimura T, Sonoda KH, Ishibashi T, Matsuda S, Ikeda E : Basigin can be a therapeutic target to restore the retinal vascular barrier function in the mouse model of diabetic retinopathy. *Sci Rep* 6, 38445, 2016 doi: 10.1038/srep38445
- 6) Ishii A, Kimura T*, Sadahiro H, Kawano H, Takubo K, Suzuki M, Ikeda E : Histological characterization of the tumorigenic "peri-necrotic niche" harboring quiescent stem-like tumor cells in glioblastoma. *PLoS ONE* 2016 Jan 22; 11 (1) : e0147366. doi: 10.1371/journal.pone.0147366. eCollection 2016
- 7) Cui D, Arima M, Takubo K, Kimura T, Horiuchi K, Minagawa T, Matsuda S, Ikeda E : ADAM12 and ADAM17 are essential molecules for hypoxia-induced impairment of neural vascular barrier function. *Sci Rep* 5 : 12796, 2015 doi : 10.1038/srep12796

- 8) Takahashi Y, Izumi Y, Kohno M, Kimura T, Kawamura M, Okada Y, Nomori H, Ikeda E : Thyroid transcription factor 1 influences the early phase of compensatory lung growth in adult mice. *Am J Respir Crit Care Med* 181 : 1397-1406, 2010
- 9) Koto T, Takubo K, Ishida S, Shinoda H, Inoue M, Tsubota K, Okada Y, Ikeda E: Hypoxia disrupts the barrier function of neural blood vessels through changes in the expression of claudin-5 in endothelial cells. *Am J Pathol* 170 : 1389-1397, 2007